

Journal Pre-proof

Management for the drowning patient

Dr David Szpilman, Dr Paddy Morgan

PII: S0012-3692(20)34896-0

DOI: <https://doi.org/10.1016/j.chest.2020.10.007>

Reference: CHEST 3690

To appear in: *CHEST*

Received Date: 27 February 2020

Revised Date: 1 October 2020

Accepted Date: 5 October 2020

Please cite this article as: Szpilman D, Morgan P, Management for the drowning patient, *CHEST* (2020), doi: <https://doi.org/10.1016/j.chest.2020.10.007>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Copyright © 2020 Published by Elsevier Inc under license from the American College of Chest Physicians.



Word Count: (also include # of references – limits 3500 words and 75 references)

Submitted for: CHEST Reviews

“Management for the drowning patient”

Dr David Szpilman – Brazil

**Brazilian Life Saving Society (SOBRASA) – Founder, Former President, Medical Director,
Medical and Prevention Commission of International Life-saving Federation (ILS) –
Member,
Science and Technology Department - Municipal Health Secretary – Rio de Janeiro City,
Drowning Resuscitation Center - GMAR – CBMERJ – Retired
<david@szpilman.com>**

Dr Paddy Morgan

**Medical Advisor, Surf Lifesaving Great Britain, UK
Medical Commission International Life Saving Federation – Member
Research Associate, Extreme Environments Laboratory, Department of Sport and Exercise
Science, University of Portsmouth.
<drpaddymorgan@gmail.com>**

Corresponding address:

David Szpilman - Av. das Américas 3555, bloco 2, sala 302, Barra da Tijuca - Rio de Janeiro – RJ - Brazil 22631-003. Phone: +55 021 999983951 david@szpilman.com

Authors has NO conflict of interest

ABSTRACT

Drowning is “*the process of experiencing respiratory impairment from submersion or immersion in liquid*”. According to WHO, drowning claim the lives of more than 40 people every hour of every day. Drowning involves some physiological principles and medical interventions that are unique. It occurs in a deceptively hostile environment that involves an underestimation of the dangers or an overestimation of water competency. It has been estimated that more than 90% are preventable. When water is aspirated into the airways, coughing is the initial reflex response. The acute lung injury alters the exchange of O₂ in different proportions. The combined effects of fluid in the lungs, loss of surfactant and increased capillary–alveolar permeability result in decreased lung compliance, increased right-to-left shunting in the lungs, atelectasis, and alveolitis, a non-cardiogenic pulmonary edema. Salt and fresh water aspiration cause similar pathology. If the person is not rescued, aspiration continues and hypoxemia leads to loss of consciousness and apnea in seconds to minutes. As a consequence, hypoxic cardiac arrest occurs. The decision to admit to an ICU should consider patient’s drowning severity and co-or-premorbidity conditions. Ventilation therapy should achieve an intrapulmonary shunt of 20% or less, or PaO₂:FiO₂ of 250 or more. Premature ventilatory weaning may cause the return of pulmonary edema with the need for re-intubation, and an anticipation of prolonged hospital stays and further morbidity. This review includes all the essentials steps from the first call to action until the best practice at the pre-hospital, emergency department, and hospitalization.

CHEST Reviews series!**MANAGEMENT FOR THE DROWNING PATIENT****IMPORTANCE OF THE TOPIC**

- Drowning is defined as the process of experiencing respiratory impairment from submersion or immersion in liquid.
- Drowning is a leading cause of injury and death among young people where it has been estimated that more than 90% are preventable.
- Mortality and morbidity are proportional to the hypoxic insult, its' treatment is the mainstay of therapy.
- Almost all drowning victims return home safely without sequelae, except the post-cardio-pulmonary arrest victims where outcome is almost solely determined by a single fate factor - duration of submersion and ICU care.
- Concurrent pathologies may "trigger" a drowning event and should be considered.

INTRODUCTION

According to the World Health Organization(WHO) drowning is a preventable public health threat claiming the lives of more than 40 people every hour of every day. With more than 90% of these deaths occurring in low- and middle-income countries, it is the world's third leading unintentional injury killer.¹ International data severely underestimates the actual drowning mortality rate in high-income countries² with survey data from some low- and middle-income countries suggesting rates four to five times that of the WHO estimated drowning rate.³ Almost all non-fatal drowning victims are able to help themselves or are rescued in time by bystanders or professional rescuers, but are rarely globally reported. Coastal drownings are estimated to cost more than \$273 million per year in the United States and more than \$228 million per year(in U.S. dollars) in Brazil.⁴ Key risk factors for drowning are male sex, age of less than 14 years, alcohol use, low income, poor education, rural residency, aquatic exposure, risky behavior and lack of supervision.^{1,4}

Drowning involves some physiological principles and medical interventions that are rarely found in other medical situations. Drowning deaths can be prevented by using a series of interventions.⁵ It occurs in a deceptively hostile environment that may not seem dangerous and usually involves an underestimation of the dangers or an overestimation of water competency to face them.⁶ The first challenge is to recognize someone at risk of drowning and appreciate

the need for rescue. Early self-rescue or rescue by others, may stop the drowning process and prevent the majority of initial and subsequent water aspiration, respiratory distress and other medical complications. The drowning process happens quickly^{7,8} but removing the victim from this environment has the potential for significant harm to the rescuer. Therefore, it's essential that all responders are aware of the complete sequence of action steps on drowning process.⁹ The details of the drowning event can assist the clinician in the hospital management of the pathophysiology that is likely to occur as a result.

DEFINITION, DATA AND DROWNING TIMELINE

*"Drowning is the process of experiencing respiratory impairment from submersion or immersion in liquid"*¹⁰. If respiratory impairment is NOT present, then this is just a rescue and not a drowning. The drowning process is a continuum, beginning with respiratory impairment as the victim's airway goes below the surface of the liquid (submersion) or when it splashes over the airways (immersion). If the victim is rescued at any time, the process of drowning is interrupted: a *non-fatal drowning*. If the victim dies at any time this is *fatal drowning*. Terms such as "near-drowning", "dry or wet drowning" and "secondary drowning" should not be used.¹⁰ A uniform way to report data for drowning resuscitation is the Utstein template for drowning resuscitation cases.¹¹ For non-fatal drownings, the WHO proposed a framework based on morbidity and severity of respiratory impairment (table 1).¹²

The drowning timeline describes every constituent of the process, triggers, actions and interventions from a temporal perspective of: Pre- event; Event and Post-event. The drowning timeline constitutes a powerful tool to improve drowning data collection, contributing to a better understanding of the process to effectively prevent, react and mitigate it and to facilitate the prioritization of cost/benefit ratios related to public health, financial aspects, political scope and social impacts.⁶ (figure 1.)

PATHOPHYSIOLOGY

Whatever the reason a person is in the water drowning carries a higher possibility of death if the individual is not rescued or unable to cope with the situation.⁶ The initial triggers for drowning are diverse and very complex.⁹ It may simply be an inability to stay afloat (e.g. young children may sink with minimal struggle¹³) or to exit the water (e.g. river channel). They vary with age, circumstance, water temperature (cold water may precipitate cardiac arrhythmias –

autonomic conflict), water competency, and some events temporally associated with being in the water e.g. Traumatic injury, or illness(myocardial infarction, seizure etc).^{14,15} These may all result in a physical inability or loss of consciousness. Many of these variables are still not fully understood. In the majority of drowning events the victim fails to keep their airway above the surface, water that enters the mouth is voluntarily spat out or swallowed. When water is aspirated into the airways, coughing occurs as an initial reflex response. Some morphological forensic studies, indicate that penetration of liquid into the lungs occurs in almost all drowning deaths. Dry-lungs can be found only in bodies disposed of into water after death on land.¹⁶ In less than 2% of cases^{17,18} laryngospasm may be present but the onset of hypoxia will terminate this rapidly. If the person is not rescued, aspiration of water continues and hypoxemia leads to loss of consciousness and apnea in seconds to minutes.^{8,19} As a consequence, hypoxic cardiac arrest generally occurs after a period of bradycardia and pulseless electrical activity and not ventricular fibrillation or tachycardia.^{20,21} Following rescue, the clinical picture is determined by the personal reactivity of the airways and the amount of water that has been aspirated with the corresponding hypoxia. Water in the alveoli causes surfactant destruction and wash-out, initiating an acute lung injury. Salt and fresh water aspiration cause similar pathology. In either situation, the effect of the osmotic gradient on the alveolar-capillary membrane can disrupt its integrity, increase its permeability and exacerbate fluid, plasma, and electrolyte shifts.²⁰ The clinical picture is of regional or generalized pulmonary edema that alters the exchange of O₂ and CO₂ in different proportions.^{19,20,22} In animal research²², the aspiration of 2.2 ml of water per kilogram of body weight lead to a severe disturbance on exchange of oxygen, decreasing the arterial oxygen pressure(PaO₂) to approximately 60 mm Hg within 3 minutes. In humans, it seems that as little as 1 to 3 ml/kg of water aspiration produces profound alterations in pulmonary gas exchange and decreases pulmonary compliance by 10% to 40%. The combined effects of fluid in the lungs, loss of surfactant and increased capillary–alveolar permeability can result in decreased lung compliance, increased right-to-left shunting in the lungs, atelectasis, and alveolitis, a non-cardiogenic pulmonary edema.²⁰

PRE-HOSPITAL CARE - DROWNING CHAIN OF SURVIVAL

The drowning chain of survival⁹(figure 2) refers to a series of interventions that, when put into action by lay or professional people, reduces the mortality associated with drowning.

- 1. Prevent drowning**-The most effective way to reduce the number of drowning deaths is prevention. It has been estimated that more than 90% of all drowning are preventable.^{5,23,24}
- 2. Recognize distress and call for help**-Recognizing a person in distress and sending for help is a key element that ensures early activation of professional rescue and medical services.⁹
- 3. Provide flotation to the victim, stop the process of drowning by reducing** the submersion risk.⁹ It is critical that personnel take precautions not to become another victim by attempting inappropriate or dangerous rescue responses.^{7,8} If not interrupted, the drowning process leads to unconsciousness and apnea, rapidly followed by cardiac arrest. During this short window of opportunity, immediate **in-water ventilation** may provide benefit if safe to do so. It can increase the discharge from hospital without sequelae by more than threefold but it is only possible if the rescuer is highly trained. Victims with only respiratory arrest usually respond after a few rescue breaths. If there is no response, the victim should be assumed to be in cardiac arrest and be rescued as quickly as possible to a location where full CPR can be initiated, as In-water chest compressions are futile.²⁵ Considering the low spinal injury incidence(0.009-0.5%), an attempt to immobilize it should only be made if there is strong indication of injury and certainly not in cases where the victim appears lifeless.^{26,27,28,29,30,31}
- 4. Remove from water - rescue only if safe to do so** - Rescue involves three phases: approach, contact and stabilization. Removal from water is essential to terminate the drowning process and allows assessment and clinical management of the victim.²⁵ Take into consideration the rescuer experience, entering the water is a personal decision. Extrication of the water is preferably in a near horizontal position, but with the head maintained above body level and airway open.³²
- 5. Provide care as needed – Basic and advanced life support**
Basic Life Support(BLS)-Cardiopulmonary or isolated respiratory arrest in drowning comprises less than 0.5% of all rescues.²³ Early basic life-support contributes to a good outcome and should be initiated as soon as possible.⁴ Once on land the victim should be placed supine, with trunk and head at the same level, and checked for responsiveness and normal breathing. If unconscious but breathing, the recovery position should be used.^{25,32} If not breathing, ventilation is essential.^{4,19,26} Hypoxia is the primary cause of cardiac arrest in drowning and requires rapid alleviation.^{8,19,26,33} Thus the Airway–Breathing–Circulation(ABC)³⁴ sequence is used, including five initial ventilations followed by 30 chest compressions. The initial 5 ventilations aims to overcome the high lung resistance, due to fluid and foam occluding the airways, allowing oxygen to reach the alveolars.^{26,35} Following this a ratio of two ventilations to

30 compressions is used until there are: signs of life; rescuer exhaustion; or advanced life support(ALS) becomes available.^{26,33} This is in preference to Circulation-Airway-Breathing(CAB) or Compression-only cardio-pulmonary resuscitation(CC-CPR) sequences^{26,33}, however any attempt at resuscitation is preferential to none. It is common for swallowed water and stomach contents to be regurgitated into the airway, with subsequent risk of aspiration.^{19,25,36} Active efforts to expel water from the airway(abdominal thrusts or placing the victim head down) should be avoided as they delay initiation of ventilations, increase the risk of vomiting by more than five-fold and thereby lead to significant increase in mortality.^{25,32} If vomiting occurs, the victims should be immediately turned onto the lateral position, vomitus removed by a finger sweep or suction and resuscitation continued. The effectiveness of automated external defibrillators (AED) for cardiac arrest in drowning is low as the presenting rhythm is usually pulse-less electrical activity or asystole. The incidence of ventricular fibrillation or ventricular tachycardia is low(4.5-6%). A shockable rhythm is however a positive predictor of survival^{8,36-38} and more likely if there is a history of coronary artery disease, epinephrine use or in the presence of severe hypothermia.²¹

Advanced Life Support (ALS) is given according to drowning severity classification^{4,19} stratified into 6 grades(algorithm 1)^{4,19} recommending the best practice treatment and the likelihood of death.

At *grade 6*(cardio-pulmonary arrest) advanced CPR should be initiated at the scene by using bag-valve-mask ventilation with high flow oxygen until a definitive airway can be achieved(oro-tracheal tube-OTT). The use of supraglottic airway devices is controversial. The pulmonary airway pressure usually exceeds safety threshold to maintain pharyngeal seals with pressures of 25–28 cm H₂O^{35,39} allowing a high potential to leak, causing new aspirations of stomach contents(water included). Once intubated, most victims can be oxygenated and ventilated effectively despite the presence of pulmonary edema in the tracheal tube. Oro-Tracheal-tube suctioning can disturb oxygenation and lung recruitment. This should be balanced against the need to ventilate and oxygenate. Peripheral venous access is a good alternative route for drug administration in the pre-hospital setting. Intraosseous access is the alternative route, if available. Endotracheal administration of drugs is not recommended in drowning.³³ Cumulative doses of epinephrine 1mg IV(or 0.01 mg/Kg) can be considered if the routine dosage fails to achieve success after the initial 5 minutes of CPR.³³ Once resuscitation attempts are successful,

an orogastric tube can be placed to reduce gastric distention and prevent further aspiration.

Recommendations for when to start and stop resuscitation are described at table 2.

4,5,8,11,19,20,21,24,25,40,41,42

Grade 5 (isolated respiratory arrest) is usually reversed by initial BLS with oxygenation and ventilation before ALS is commenced. If there is spontaneous ventilation but oxygenation is compromised (acute pulmonary edema (*grade 3 and 4*)) the objective is to achieve a pre-hospital peripheral saturation above 92% by administering oxygen by face mask at a rate of 15 liters of oxygen/min. Early oral tracheal intubation (OTI) and mechanical ventilation are indicated as soon as possible, because of respiratory fatigue, despite adequate oxygenation by face mask. While all grade 4 need OTI, a few grades 3 drowning cases will tolerate non-invasive ventilatory (NIV) support^{19,40,43,44} provided their conscious level allows. Patients should be anaesthetized to tolerate intubation and artificial mechanical ventilation. Emergency department (ED) attendance is recommended for all grade 2 to 6 patients. Most victims *grade 2* require low flow oxygen and will normalize their clinical situation within 6–48 hours and can be discharged home.¹⁹

HOSPITAL CARE

The decision to admit to an ICU or hospital bed versus observation in the emergency department or discharge home should consider patient's drowning severity and co-or-premorbidity conditions. A thorough medical assessment should be performed including, chest radiography and/or lung ultrasound, arterial blood gas measurement and blood tests.^{45,46} The latter should include tests of renal function, liver function, electrolytes, hemoglobin and any appropriate toxicology given the association with suicide and alcohol excess. Patient grade 3 to 6 should be admitted to an Intensive Care Unit (ICU) for close observation and therapy. Patient grade 2 can be observed in the emergency room, but grade 1 and rescue cases with no complaints or associated illness or trauma can be released home.^{19,45}

Respiratory system

Patient grade 3 to 6 usually will arrive from pre-hospital ALS on mechanical ventilation with acceptable oxygenation. If not, the physician should reassure that. Oxygen start at 100%, but should be reduced as soon as possible. Positive end-expiratory pressure (PEEP) should be added initially at a level of 5 cm H₂O and then increased by 2 to 3 cm H₂O increments if needed and possible. The PEEP should be used until the desired intrapulmonary shunt (QS: QT) of 20% or less, or PaO₂:FiO₂ of 250 or more is achieved.²⁰ At *grade 4* if hypotension is not corrected by

oxygen, a rapid crystalloid infusion should be used before trying to reduce PEEP. Once the desired oxygenation is achieved, that level of PEEP should be maintained unchanged for at least 48 hours before attempting to weaning.⁴⁷ This is the minimum time required for adequate surfactant regeneration. Premature ventilatory weaning may cause the return of pulmonary edema with the need for re-intubation, and an anticipation of prolonged hospital stays and further morbidity.⁴ A clinical picture very similar to acute respiratory distress syndrome (ARDS), but with a prompt recovery and no lung sequelae is common after significant drowning episodes (grade 3 to 6). A protective lung ventilation strategy (e.g.; low tidal volumes [6 mL/kg ideal body weight]) similar to ARDS should be used. Permissive hypercapnia should be avoided however to prevent further neurological insult in those with significant hypoxic-ischemic brain injury (usually grade 6).⁴ Continuous Positive Airway Pressure (CPAP), Pressure Support Ventilation mode (PSV) and/or NIV are appropriate weaning strategies if pulmonary and psychological status allows.^{43,44}

Pools, rivers and beaches generally have insufficient bacteria colonization to promote pneumonia in the immediate post drowning period⁴⁸. Pneumonia is often misdiagnosed initially because of the early radiographic appearance of water in the lungs with few actually requiring antibiotic therapy (12%). If the victim requires mechanical ventilation, the incidence of pneumonia (ventilator-associated) increases to 34-52% in the third or fourth day of hospitalization when pulmonary edema is resolving.⁴⁹ Vigilance not only for pulmonary but also other infectious complications is important. Prophylactic antibiotics tend to only select out more resistant and aggressive organisms.⁵⁰ The first signs of pulmonary infection are at 48 to 72 hours and are gauged by prolonged fever, sustained leukocytosis, persistent or new pulmonary infiltrates, and leukocyte response in the tracheal aspirates. A broad-spectrum antibiotic therapy to cover gram-positive and gram-negative should be used immediately if the drowning occurred in water with high pathogen load (UFC > 10²⁰). In ventilator associated pneumonia, the predominant microorganisms of the ICU or available cultures should be considered. In resistant infections, consideration should be given to alternate pathogens e.g. fungal, algae and protozoa. Fiberoptic bronchoscopy may be useful for evaluation of infection by obtaining quantitative cultures, determining the extent and severity of airway injury and for the rare occasions where therapeutic clearing of sand, gravel or other solids is indicated. Corticosteroids should not be used except for bronchospasm. The clinician must be aware of and constantly vigilant for volutrauma and barotrauma during mechanical ventilation.⁴⁸

Focused bedside ultrasound can be used to diagnose and monitor the respiratory and circulatory system in real time. Specifically, the rapid diagnosis of pneumothoraces and distribution of lung oedema.⁴⁶ Ultrasound assessment of cardiac function can guide fluid therapy, indication for inotropes or vasopressors, monitor the response to therapy and exclude concurrent pathologies.

Circulatory system

Cardiac dysfunction with low cardiac output is usual immediately after severe cases.²⁰ Low cardiac output is associated with high pulmonary capillary occlusion pressure (hypoxic vasoconstriction), high central venous pressure and pulmonary vascular resistance which can persist for days. This may add a cardiogenic component to the drowning primary non-cardiogenic pulmonary edema. The reduced cardiac output can be corrected with oxygenation, crystalloid infusion and restoration of normal body temperature. Vasopressor infusion should be reserved for refractory hypotension. Echocardiography to assess cardiac function can guide the clinician in titrating inotropes, vasopressors or both if volume crystalloid replacement is inadequate⁴. In patients who are hemodynamically unstable or have severe pulmonary dysfunction, pulmonary artery catheterization may be considered to provide useful information. There is no evidence to support the use of any specific fluid therapy for salt and fresh water drowning²⁰, the use of diuretics or water restriction. Metabolic acidosis occurs in 70% of patients arriving at the hospital after a drowning episode.¹⁹

Neurologic system

Most late deaths and long-term sequelae of drowning are neurologic in origin (anoxic-ischemic cerebral insult) and are almost exclusive in grade 6, as pulmonary injury is usual reversible.^{4,48} Although the highest priority of CPR is restoration of spontaneous circulation, every effort in the early stages should be directed at resuscitating the brain and preventing further neurologic damage. These steps include providing adequate oxygenation (SatO₂p > 92%) and cerebral perfusion (mean arterial pressure around 100 mm Hg). Any victim who remains comatose or unresponsive after successful CPR or deteriorates neurologically should undergo careful and frequent neurologic function assessment and care by using the measures at table 3.^{40,51}

Drowning victims with spontaneous circulation who remain comatose should have targeted temperature management to improve outcomes after cerebral hypoxia-ischaemia.⁵²⁻⁵⁴ Maintaining a core temperature of 32°C to 34°C for at least 24 hours post arrest is associated with improved neurological outcomes.⁵²⁻⁵⁵ Although there is insufficient evidence to support a specific target PaCO₂ or oxygen saturation, hypoxemia should be avoided. Studies have failed to demonstrate improved outcome utilizing intracranial pressure monitoring⁵⁶, therapies to control intracranial hypertension or maintenance of artificially high cerebral perfusion pressure(CPP).^{56,57}

New therapeutic interventions for drowning victims such as artificial surfactant⁵⁸⁻⁶⁰ or nitric oxide⁶¹ are still experimental with a few successful case reports. Extracorporeal membrane oxygenation may be considered when the patient is profoundly hypothermic, or conventional respiratory assistance is insufficient to maintain oxygenation. This assumes it is available and feasible.^{62,63,64,65}

Unusual complications

In the most severe cases(grade 6) the hypoxic and/or hypo-perfusion associated with drowning can trigger the systemic inflammatory response syndrome. This can manifest as isolated cardiac, renal or hepatic dysfunction through to sepsis and multi-organ dysfunction syndrome. Rarely, drowning victims with normal chest radiography develop fulminant pulmonary edema upto 12 hours after the incident. Whether this late-onset pulmonary edema is delayed ARDS, a neurogenic pulmonary edema secondary to hypoxia, or just an airway hyperreactive to water aspiration is still unclear.^{4,19,11,66}

Cold water drowning

Submersion in cold water and sudden release of breath hold can induce cardiac arrhythmias, particular in those with long QT syndrome, by simultaneously activating the antagonistic responses of the autonomic nervous system – autonomic conflict. These being the sympathetic “cold shock response” producing a tachycardia and the parasympathetic “diving response” mediated bradycardia. This may, in vulnerable individuals, account for sudden death in cold water.⁵⁷⁻⁶⁹ Hypothermia reduces the electrical and metabolic activity of the brain; with cerebral oxygen consumption reducing by approximately 5% per each °C reduction in temperature within the range of 37 to 20°C.⁷⁰ Thus, prolonging the interval until cellular anoxia, ATP depletion and

cell death signaling. If cooling occurs prior to submersion (i.e. hypoxia) it provides a form of cerebral protection that explains cases with good neurological outcome despite prolonged submersion for up to 90 minutes.^{26,41} In these cases, the water temperature was 6°C or lower. Extracorporeal membrane oxygenation(ECMO) has been used in the resuscitation of these victims with good neurological outcomes, despite initially poor prognosis.^{4,33,41,62,63} Previously hypothermic victims re-warmed to near-normal core temperature, who remain asystolic with a significantly elevated serum potassium, despite resuscitation, is a key indicator of futility.^{26,71,72}

OUTCOME AND SCORING SYSTEMS

Of the drowning grades 1 to 5, 95% return home without sequelae.¹⁹ In grade 6, prognostic variables are important while counseling family members and in deciding which treatment strategies are appropriate.⁸ Victims who remain comatose or deteriorate neurologically should undergo intensive assessment and care.⁴⁰ Several studies have established that outcome is almost solely determined by a single fate factor-duration of submersion(table 4).^{8,11,19,20,21,24,25,40,41,42}

Error! Reference source not found.This emphasizes the need for accurate documentation of the pre-hospital presentation and incident details. After successful CPR, assessment of neurological severity will allow comparison of different therapeutic approaches.⁷³ Data suggest that patients who remain decorticate, decerebrate, or flaccid in the 2 to 6 hours after the drowning incident (when no drugs are implicated) are brain dead or will survive with moderate to severe neurological impairment.⁵ Patients who are improving but remain unresponsive have a 50% likelihood of a good outcome(table 5).⁸

Conclusion

As one of the most common causes of unintentional injury-related morbidity and mortality worldwide, drowning remains a significant public health issue and an extremely complex process in which there is no simple, or single solution. However, the true impact of drowning on public health is unknown due to a lack of high-quality epidemiological data in the field. The most effective intervention to reduce drowning deaths is prevention. When prevention fails further reduction in morbidity and mortality is only achieved by effective rescue, and early clinical interventions when indicated. In many areas of medicine, it is obvious that prevention is better than cure, but how do you motivate and educate those populations that are at the highest risk? Does it require the emotive scenario of a child death or severe neurological insult as a result of a drowning event for people to act? The drowning process may involve a complex interplay between acute injury or disease and an inability to maintain the airway clear of the waters surface. The simple life skills of water awareness and the ability to float face up will prevent many of the complications of this potentially fatal process. There is a deficit of high-quality scientific evidence at all stages of the patient's journey following a drowning event, particularly in the hospital setting. These events result in a multisystem disorder to a greater or lesser extent depending upon the duration of the hypoxic insult. Following a successful rescue, the key therapy is oxygen, and facilitation of its' delivery to the tissues of the body.

REFERENCES

1. Bloomberg LP, World Health Organization, editors. Global report on drowning: preventing a leading killer. Geneva, Switzerland: World Health Organization; 2014.
2. Lu T-H, Lunetta P, Walker S. Quality of cause-of-death reporting using ICD-10 drowning codes: a descriptive study of 69 countries. *BMC Med Res Methodol* 2010;10:30.
3. Peden M, Organization WHO, others. World report on child injury prevention. Geneva: World Health Organization; 2008.
4. Szpilman D, Bierens JJLM, Handley AJ, Orłowski JP. Drowning. *N Engl J Med* 2012;366(22):2102–2110.
5. Moran K, Quan L, Franklin R, Bennett E. Where the Evidence and Expert Opinion Meet: A Review of Open-Water Recreational Safety Messages. *Int J Aquat Res Educ* [Internet] 2011;5(3). Available from: <https://scholarworks.bgsu.edu/ijare/vol5/iss3/5>
6. Szpilman D, Tipton M, Sempsrott J, et al. Drowning timeline: a new systematic model of the drowning process. *Am J Emerg Med* 2016;34(11):2224–2226.
7. Venema AM, Groothoff JW, Bierens JJLM. The role of bystanders during rescue and resuscitation of drowning victims. *Resuscitation* 2010;81(4):434–439.

8. Orlowski JP, Szpilman D. Drowning. Rescue, resuscitation, and reanimation. *Pediatr Clin North Am* 2001;48(3):627–646.
9. Szpilman D, Webber J, Quan L, et al. Creating a drowning chain of survival. *Resuscitation* 2014;85(9):1149–1152.
10. Beeck EF van, Branche CM, Szpilman D, Modell JH, Bierens JJLM. A new definition of drowning: towards documentation and prevention of a global public health problem. *Bull World Health Organ* 2005;83(11):853–856.
11. Idris AH, Bierens JJLM, Perkins GD, et al. 2015 Revised Utstein-Style Recommended Guidelines for Uniform Reporting of Data From Drowning-Related Resuscitation: An ILCOR Advisory Statement. *Circ Cardiovasc Qual Outcomes* 2017;10(7).
12. WHO | Non-fatal Drowning Terminology [Internet]. WHO. [cited 2020 Jan 5]; Available from: http://www.who.int/violence_injury_prevention/drowning/non-fatal-drowning/en/
13. Langendorfer S. Applying a developmental perspective to aquatics and swimming. *Biomech Med Swim XI* 2010;20–22.
14. Szpilman D, Orlowski JP. Sports related to drowning. Number 2 in the Series “Sports-related lung disease”. *Eur Respir Rev* 2016; 25: 348–359
15. Joost J. L. M. Bierens JJLM, Lunetta P, Tipton M, Warner DS. Physiology of drowning: A review. *PHYSIOLOGY* 31: 147–166, 2016.
16. Lunetta P, Modell JH, Sajantila A. What is the incidence and significance of “dry-lungs” in bodies found in water? *Am J Forensic Med Pathol* 25: 291–301, 2004.
17. Lunetta P, Modell JH, Sajantila A. What is the incidence and significance of “dry-lungs” in bodies found in water? *Am J Forensic Med Pathol* 2004;25(4):291–301.
18. Szpilman D, Elmann J, Cruz-Filho FE. Dry-drowning-fact or myth. In: World Congress on Drowning, Netherlands. 2002.
19. Szpilman D. Near-drowning and drowning classification: a proposal to stratify mortality based on the analysis of 1,831 cases. *Chest* 1997;112(3):660–665.
20. Orlowski JP, Abulleil MM, Phillips JM. The hemodynamic and cardiovascular effects of near-drowning in hypotonic, isotonic, or hypertonic solutions. *Ann Emerg Med* 1989;18(10):1044–1049.
21. Grmec S, Strnad M, Podgorsek D. Comparison of the characteristics and outcome among patients suffering from out-of-hospital primary cardiac arrest and drowning victims in cardiac arrest. *Int J Emerg Med* 2009;2(1):7–12.
22. Modell JH, Moya F, Newby EJ, Ruiz BC, Showers AV. The effects of fluid volume in seawater drowning. *Ann Intern Med* 1967;67(1):68–80.

23. Szpilman D, Barros Oliveira R de, Mocellin O, Webber J. Is drowning a mere matter of resuscitation? *Resuscitation* 2018;129:103–106.
24. Quan L, Pilkey D, Gomez A, Bennett E. Analysis of paediatric drowning deaths in Washington State using the child death review (CDR) for surveillance: what CDR does and does not tell us about lethal drowning injury. *Inj Prev* 2011;17(Suppl I):i28–i33.
25. Szpilman D, Soares M. In-water resuscitation-is it worthwhile? *Resuscitation* 2004;63(1):25–31.
26. Truhlář A, Deakin CD, Soar J, et al. European Resuscitation Council Guidelines for Resuscitation 2015: Section 4. Cardiac arrest in special circumstances. *Resuscitation* 2015;95:148–201.
27. Wernicki P, Szpilman D. Immobilization and extraction of spinal injuries. In: *Drowning*. Springer; 2014. p. 621–628.
28. Turgut A, Turgut T. A study on rescuer drowning and multiple drowning incidents. *J Safety Res* 2012;43(2):129–132.
29. Szpilman D, Brewster C, Cruz-Filho FES. Aquatic Cervical Spine Injury – How often do we have to worry? World Congress on Drowning. Netherlands 2002, Oral Presentation. DOI: 10.13140/2.1.4504.2886.
30. Szpilman D. Aquatic cervical and head trauma: nobody told me it could be a jump in the darkness! Danang, Vietnam: 2011. p. 153.
31. Watson RS, Cummings P, Quan L, Bratton S, Weiss NS. Cervical spine injuries among submersion victims. *J Trauma* 2001;51(4):658–662.
32. Szpilman D, Handley A. Positioning the drowning victim. In: *Handbook on drowning: Prevention, rescue, treatment*. Berlin: Springer Science & Business Media; 2006. p. 336–341.
33. Vanden Hoek TL, Morrison LJ, Shuster M, et al. Part 12: cardiac arrest in special situations: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2010;122(18 Suppl 3):S829-861.
34. Kitamura T, Iwami T, Kawamura T, et al. Conventional and chest-compression-only cardiopulmonary resuscitation by bystanders for children who have out-of-hospital cardiac arrests: a prospective, nationwide, population-based cohort study. *Lancet Lond Engl* 2010;375(9723):1347–1354.
35. Baker PA, Webber JB. Failure to ventilate with supraglottic airways after drowning. *Anaesth Intensive Care* 2011;39(4):675–677.
36. Manolios N, Mackie I. Drowning and near-drowning on Australian beaches patrolled by life-savers: a 10-year study, 1973-1983. *Med J Aust* 1988;148(4):165–167, 170–171.

37. Dyson K, Morgans A, Bray J, Matthews B, Smith K. Drowning related out-of-hospital cardiac arrests: Characteristics and outcomes. *Resuscitation* 2013;84(8):1114–1118.
38. Claesson A, Lindqvist J, Herlitz J. Cardiac arrest due to drowning—Changes over time and factors of importance for survival. *Resuscitation* 2014;85(5):644–648.
39. Benger JR, Kirby K, Black S, et al. Effect of a Strategy of a Supraglottic Airway Device vs Tracheal Intubation During Out-of-Hospital Cardiac Arrest on Functional Outcome: The AIRWAYS-2 Randomized Clinical Trial. *JAMA* 2018;320(8):779–791.
40. Topjian AA, Berg RA, Bierens JJLM, et al. Brain Resuscitation in the Drowning Victim. *Neurocrit Care* 2012;17(3):441–467.
41. Tipton MJ, Golden FSC. A proposed decision-making guide for the search, rescue and resuscitation of submersion (head under) victims based on expert opinion. *Resuscitation* 2011;82(7):819–824.
42. Quan L, Bierens JJLM, Lis R, Rowhani-Rahbar A, Morley P, Perkins GD. Predicting outcome of drowning at the scene: A systematic review and meta-analyses. *Resuscitation* 2016;104:63–75.
43. Kim JH, Sun KH, Park YJ. The Utility of Non-Invasive Nasal Positive Pressure Ventilation for Acute Respiratory Distress Syndrome in Near Drowning Patients. *J Trauma Inj* 2019;32(3):136–142.
44. Çağlar A, Er A, Özden Ö, et al. Efficacy of Early Noninvasive Ventilation in Three Cases of Nonfatal Drowning with Pulmonary Oedema in the Paediatric Emergency Department. *Hong Kong J Emerg Med* 2016;23(2):42–46.
45. Szpilman D, Semprcott J, Schmidt A. Drowning. *BMJ Best Practice*. Nov 2017. <http://bestpractice.bmj.com/topics/en-gb/657>. Last accessed 20 April 2020.
46. Lichtenstein, DA. Current Misconceptions in Lung Ultrasound - A Short Guide for Experts. *CHEST* 2019; 156(1):21-25
47. Michelet, P., et al. (2017). "Acute respiratory failure after drowning: a retrospective multicenter survey." *European Journal of Emergency Medicine* 24(4): 295.
48. Orłowski JP. Drowning, near-drowning, and ice-water submersions. *Pediatr Clin North Am* 1987;34(1):75–92.
49. Berkel M van, Bierens JJ, Lie RL, et al. Pulmonary oedema, pneumonia and mortality in submersion victims; a retrospective study in 125 patients. *Intensive Care Med* 1996;22(2):101–107.
50. Wood C. Towards evidence-based emergency medicine: best BETs from the Manchester Royal Infirmary. BET 1: prophylactic antibiotics in near-drowning. *Emerg Med J EMJ* 2010;27(5):393–394.
51. NICE-SUGAR Study Investigators, Finfer S, Chittock DR, et al. Intensive versus conventional glucose control in critically ill patients. *N Engl J Med* 2009;360(13):1283–1297.

52. Nolan JP, Soar J, Cariou A, et al. European Resuscitation Council and European Society of Intensive Care Medicine Guidelines for Post-resuscitation Care 2015: Section 5 of the European Resuscitation Council Guidelines for Resuscitation 2015. *Resuscitation* 2015;95:202–222.
53. Gunn AJ, Thoresen M. Hypothermic Neuroprotection. *NeuroRX* 2006;3(2):154–169.
54. Froehler MT, Geocadin RG. Hypothermia for neuroprotection after cardiac arrest: Mechanisms, clinical trials and patient care. *J Neurol Sci* 2007;261(1):118–126.
55. Abdul Aziz KA, Meduoye A. Is pH-stat or alpha-stat the best technique to follow in patients undergoing deep hypothermic circulatory arrest? *Interact Cardiovasc Thorac Surg* 2010;10(2):271–282.
56. Bohn D, Biggar W, Smith C, Conn A, Barker G. Influence of hypothermia, barbiturate therapy, and intracranial pressure monitoring on morbidity and mortality after near-drowning. *Crit Care Med* 1986;14(6):529–534.
57. Donnino Michael W., Andersen Lars W., Berg Katherine M., et al. Temperature Management After Cardiac Arrest. *Circulation* 2015;132(25):2448–2456.
58. Varisco BM, Palmatier CM, Alten JA. Reversal of intractable hypoxemia with exogenous surfactant (calfactant) facilitating complete neurological recovery in a pediatric drowning victim. *Pediatr Emerg Care* 2010;26(8):571–573.
59. Cubattoli L, Franchi F, Coratti G. Surfactant therapy for acute respiratory failure after drowning: Two children victim of cardiac arrest. *Resuscitation* 2009;80(9):1088–1089.
60. Fettah N, Dilli D, Beken S, Zenciroglu A, Okumuş N. Surfactant for acute respiratory distress syndrome caused by near drowning in a newborn. *Pediatr Emerg Care* 2014;30(3):180–181.
61. Takano Y, Hirosako S, Yamaguchi T, et al. [Nitric oxide inhalation as an effective therapy for acute respiratory distress syndrome due to near-drowning: a case report]. *Nihon Kokyuki Gakkai Zasshi* 1999;37(12):997–1002.
62. Wang C-H, Chou C-C, Ko W-J, Lee Y-C. Rescue a drowning patient by prolonged extracorporeal membrane oxygenation support for 117 days. *Am J Emerg Med* 2010;28(6):750.e5–7.
63. Guenther U, Varelmann D, Putensen C, Wrigge H. Extended therapeutic hypothermia for several days during extracorporeal membrane-oxygenation after drowning and cardiac arrest Two cases of survival with no neurological sequelae. *Resuscitation* 2009;80(3):379–381.
64. Griffiths MJD, et al. (2019). "Guidelines on the management of acute respiratory distress syndrome." *BMJ Open Respir Res* 6(1): e000420.
65. Svendsen, O. S., et al. (2017). "Outcome After Rewarming From Accidental Hypothermia by Use of Extracorporeal Circulation." *Annals of Thoracic Surgery* 103(3): 920.

66. Dyson, K., et al. (2013). "Drowning related out-of-hospital cardiac arrests: characteristics and outcomes." *Resuscitation* 84(8): 1114.
67. Shattock MJ, Tipton MJ. "Autonomic conflict": a different way to die during cold water immersion? *J Physiol* 2012;590(14):3219–3230.
68. Winter J, Tipton MJ, Shattock MJ. Autonomic conflict exacerbates long QT associated ventricular arrhythmias. *J Mol Cell Cardiol* 2018;116:145–154.
69. Vincenzi FF. Drug-induced long QT syndrome increases the risk of drowning. *Med Hypotheses* 2016;87:11–13.
70. Polderman KH. Application of therapeutic hypothermia in the ICU: opportunities and pitfalls of a promising treatment modality. Part 1: Indications and evidence. *Intensive Care Med* 2004;30(4):556–575.
71. Paal P, Gordon L, Strapazzon G, et al. Accidental hypothermia—an update. *Scand J Trauma Resusc Emerg Med* 2016;24(1):111.
72. Brugger H, Bouzat P, Pasquier M, et al. Cut-off values of serum potassium and core temperature at hospital admission for extracorporeal rewarming of avalanche victims in cardiac arrest: A retrospective multi-centre study. *Resuscitation* 2019;139:222–229.
73. Modell JH, Conn AW. [Current neurological considerations in near-drowning]. *Can Anaesth Soc J* 1980;27(3):197–198.

TABLES, FIGURES AND ALGORITHM**Table 1 - Categorization Framework for Non-fatal Drowning¹²**

1. The severity of respiratory impairment immediately after the drowning process stopped. There must be evidence of respiratory impairment to be classified as a non-fatal drowning.
2. The morbidity category at the time when non-fatal drowning information is gathered. For the purposes of this categorization framework, morbidity is defined as a decline from the individual's functional capacity prior to the drowning.

^aThe following descriptors serve to better characterize the meaning of "involuntary distressed coughing": coughing up liquid / moving liquid out of the airway; in water, in distress and coughing; sustained coughing. ^bThe phrase "previous functional capacity" includes the person's cognitive, motor, and psychological capacity (WHO task force – personal communication)

(attached in word file)

Figure 1 – drowning timeline⁶**Figure 2 – Drowning Chain of Survival⁹**

ALGORITHM 1 - Drowning severity classification and flow chart strategy decision based on evaluation of 87,339 rescues Error! Reference source not found.,Error! Reference source not found.

Table 2 – Drowning – When to initiate CPR and when to discontinue^{4,5,8,11,19,20,21,24,25,40,41,42}

Table 3 – Recommended care for victim who remains comatose or unresponsive after successful CPR or deteriorates neurologically^{40,51}

Table 4 - Probability of Neurologically Intact Survival to Hospital Discharge, based on duration of submersion^{8,11,19,20,21,24,25,40,41,42}

Table 5 – Clinical Prognostic Score for the immediate period pos successful CPR, based on Glasgow Coma Score.⁸

Severity of respiratory impairment after the drowning process stopped.		
(1) Mild	(2) Moderate	(3) Severe
<ul style="list-style-type: none"> Breathing Involuntary distressed coughing^a AND Fully alert 	<ul style="list-style-type: none"> Difficulty breathing AND Disoriented but conscious 	<ul style="list-style-type: none"> Not breathing AND Unconscious
Morbidity category (based upon any decline from previous functional capacity ^b) at the time of measurement.		
(A) No morbidity	(B) Some morbidity	(C) Severe morbidity
<ul style="list-style-type: none"> No decline 	<ul style="list-style-type: none"> Some decline 	<ul style="list-style-type: none"> Severe decline

Table 1 - Categorization Framework for Non-fatal Drowning¹²

1. The severity of respiratory impairment immediately after the drowning process stopped. There must be evidence of respiratory impairment to be classified as a non-fatal drowning.
2. The morbidity category at the time when non-fatal drowning information is gathered. For the purposes of this categorization framework, morbidity is defined as a decline from the individual's functional capacity prior to the drowning.

^aThe following descriptors serve to better characterize the meaning of "involuntary distressed coughing": coughing up liquid / moving liquid out of the airway; in water, in distress and coughing; sustained coughing. ^bThe phrase "previous functional capacity" includes the person's cognitive, motor, and psychological capacity (WHO task force – personal communication)

Question	Recommendations
In whom to begin?	<ul style="list-style-type: none"> • Give ventilatory support for respiratory distress/arrest to avoid cardiac arrest. • Start CPR in all submersions < 60 minutes who does not present obvious physical evidence of death (rigor mortis, body decomposition or dependent lividity).
When to discontinue?	<ul style="list-style-type: none"> • Basic life support should continue unless signs of life re-appear, rescuers exhaustion or advanced life support take over. • Advanced life support should be ongoing until patient has been rewarmed (if hypothermic) and asystole persist for more than 20 minutes.

Table 2 – Drowning – When to initiate CPR and when to discontinue^{4,5,8,11,19,20,21,24,25,40,41,42}

Journal Pre-proof

Raise the head of the bed by 30 degrees (if there is no hypotension);
Maintain adequate mechanical ventilation by using drugs to low patient fighting the ventilator;
Ensure appropriate respiratory toilet (keeping positive airway pressure) without provoking hypoxia;
Treat for seizure activity;
Avoid sudden metabolic corrections;
Prevent interventions that increase intracranial pressure (ICP) - including urinary retention, pain, hypotension, hypercapnia, hypoxemia;
Hyperthermia should be avoided and normoglycemia maintained.

Table 3 – Recommended care for victim who remains comatose or unresponsive after successful CPR or deteriorates neurologically^{40,51}

Journal Pre-proof

Duration of submersion	Death or severe neurological impairment
0 to <5 minutes	10%
5 to <10 minutes	56%
10 to <25 minutes	88%
> 25 minutes	99,9%

Note in these data how 5 more minutes of submersion in the 5 to <10 min group increases mortality almost 6 times compared to the 0 to <5-minute group


Table 4 - Probability of neurologically intact survival to hospital discharge, based on duration of submersion^{8,11,19,20,21,24,25,40,41,42}

NEUROLOGIC PROGNOSTIC SCORE	
Post successful CPR on Drowning	
A – FIRST HOUR	B – AFTER 5 to 8 h
Alert - 10	Alert - 9.5
Confused - 9	Confused - 8
Torpor - 7	Torpor - 6
Coma with normal brainstem - 5	Coma with normal brainstem - 3
Coma with abnormal brainstem - 2	Coma with abnormal brainstem - 1
A + B	
RECOVERY WITHOUT SEQUALAE	
Excellent (≥ 13)	$\geq 95\%$
Very good (10-12)	75 to 85%
Good (8)	40 to 60%
Regular (5)	10 to 30%
Poor (3)	$\leq 5\%$

Table 5 – Drowning clinical prognostic score for the immediate period pos successful CPR, based on Glasgow Coma Score.⁸

DROWNING TIMELINE


SYSTEMATIC MODEL OF THE DROWNING PROCESS



Szpilman D, Tipton M, Sempstrott J, Webber J, Bierens J, Dawes P, Seabra R, Barcala-Furelos R, Queiroga AC. Drowning timeline: a new systematic model of the drowning process, Am J Emerg Med. 2016 Nov;34(11):2224-2226. Authors acknowledge SEMES (Sociedad Espanola de Medicina de Urgencias e Emergencias) for the design.

DROWNING CHAIN OF SURVIVAL

A call to action



Journal Pre-proof